

Polygynax

This information is a summary only. It does not contain all information about this medicine. If you would like more information about the medicine you are taking, check with your doctor or other health care provider. No rights can be derived from the information provided in this medicine leaflet.

Product

Polygynax® vaginal capsules

Composition:

Active ingredients : Neomycin sulphate 35,000 IU • Polymyxin B sulphate 35,000 IU • Nystatin 100,000 IU • Excipients: Tefose 63®, hydrogenised soybean oil, dimethylpolysiloxan, gelatin and glycerol. Boxes of 6 and 12 vaginal capsules.

Therapeutic category:

Topical anti-infectious agent (G. genitourinary tract).

Indication:

Topical treatment of vaginal infections.

Contra indications:

Allergy to one of its ingredients or another substance of the same family

- use of latex diaphragms or condoms. Inform your physician or your pharmacist about all medication you are using with Polygynax® at the same time.

Special Warnings:

Polygynax® contains soybean oil. If you are allergic to peanut or soya, do not use this medicinal product. Stop treatment in case of local intolerance or allergic reactions. In case of local allergy, the subsequent use of the same antibiotic or related antibiotics via the oral route may be compromised.

Precautions:

Inform your physician in case of renal failure. Do not exceed the treatment duration recommended by your physician.

Interactions:

Do not use with spermicidal products (risk of inactivation) and condoms

Adverse effects:

Possibility of contact allergic eczema or away from the site of application in case of prolonged use. Report any unwanted and bothersome effects not mentioned in this insert to your physician or pharmacist.

Pregnancy Lactation:

the use of this drug is not recommended during pregnancy. This drug must not be taken while breast-feeding. In general, pregnant women and nursing mothers should always ask their physician or pharmacist for advice before taking any drugs.

Dosage - Reserved for Adults:

Place one capsule deep in the vagina for 12 days. It is preferable and easier to insert the capsule while lying down on your back with the knees apart.

Practical advice :

- to avoid contaminating the people around you, use your own face cloth and towel
- wear cotton underwear
- avoid vaginal douches
- avoid using tampons during treatment
- do not stop treatment during menstrual periods
- your physician may also treat your sexual partner in order to avoid reinfection.

Storage Conditions:

Use before the expiry date indicated on the external packaging. Store below 25°C.

DO NOT SWALLOW. KEEP OUT OF THE REACH AND SIGHT OF CHILDREN.